



Early Journal Content on JSTOR, Free to Anyone in the World

This article is one of nearly 500,000 scholarly works digitized and made freely available to everyone in the world by JSTOR.

Known as the Early Journal Content, this set of works include research articles, news, letters, and other writings published in more than 200 of the oldest leading academic journals. The works date from the mid-seventeenth to the early twentieth centuries.

We encourage people to read and share the Early Journal Content openly and to tell others that this resource exists. People may post this content online or redistribute in any way for non-commercial purposes.

Read more about Early Journal Content at <http://about.jstor.org/participate-jstor/individuals/early-journal-content>.

JSTOR is a digital library of academic journals, books, and primary source objects. JSTOR helps people discover, use, and build upon a wide range of content through a powerful research and teaching platform, and preserves this content for future generations. JSTOR is part of ITHAKA, a not-for-profit organization that also includes Ithaka S+R and Portico. For more information about JSTOR, please contact support@jstor.org.

446. Proposed by S. G. BARTON, University of Pennsylvania.

Prove any one or more of the sixteen theorems, stated without proof, in the article in this issue of the MONTHLY (pages 182-184) on "Properties of the Normals to a Conic."

CALCULUS.

366. Proposed by I. A. BARNETT, University of Chicago.

Compute the definite integral $\int_a^b \sin^{-1} x dx$ where $0 \leq a \leq 1$ and $0 \leq b \leq 1$, by direct summation.

367. Proposed by C. N. SCHMALL, New York City.

Show that the volume enclosed by the surface $(x^2 + y^2 + z^2)^5 = (a^2x^2 + b^2y^2 + c^2z^2)^2$ is $\frac{4}{3}\pi(a^3 + b^3 + c^3)$.

MECHANICS.

295. Proposed by B. F. FINKEL, Drury College.

A homogeneous hollow cylinder, whose inner radius is half of its outer radius, rolls without slipping down a plane inclined at an angle α to the horizontal. Find its acceleration.

[From Prescott's *Mechanics of Particles and of Rigid Bodies*.]

NUMBER THEORY.

218. Proposed by ELIJAH SWIFT, Princeton, N. J.

If p is prime and > 3 , show that $\sum_{a=1}^{a=p-1} 1/a^2 \equiv 0 \pmod{p}$.

219. Proposed by R. D. CARMICHAEL, Indiana University.

Determine whether it is possible for a polygon to have the number of its diagonals equal to a perfect fourth power.

SOLUTIONS OF PROBLEMS.

ALGEBRA.

A solution of 399 by WM. CULLUM and a solution of 400 by LOUIS O'SHAUGHNESSY were received too late for credit in the May issue.

401. Proposed by R. D. CARMICHAEL, Indiana University.

Prove the validity of Borda's series:

$$\log(x+2) = 2 \log(x+1) - 2 \log(x-1) + \log(x-2) \\ + 2 \left[\frac{2}{x^3-3x} + \frac{1}{3} \left(\frac{2}{x^3-3x} \right)^3 + \frac{1}{5} \left(\frac{2}{x^3-3x} \right)^5 + \cdots \right].$$

SOLUTION BY A. M. HARDING, University of Arkansas.

We have

$$\log(x+2) - 2 \log(x+1) + 2 \log(x-1) - \log(x-2) \\ = \log \frac{(x-1)^2(x+2)}{(x+1)^2(x-2)} = \log \frac{x^3-3x+2}{x^3-3x-2} = \log \frac{1 + \frac{2}{x^3-3x}}{1 - \frac{2}{x^3-3x}} \\ = 2 \left[\frac{2}{x^3-3x} + \frac{1}{3} \left(\frac{2}{x^3-3x} \right)^3 + \frac{1}{5} \left(\frac{2}{x^3-3x} \right)^5 + \cdots \right],$$

which proves the result.